

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE October 2000	3. REPORT TYPE AND DATES COVERED Final Report: Dec. 1991 - Nov. 1998		
4. TITLE AND SUBTITLE Electron-Molecule Collisions and UV Signatures in Flowfields		5. FUNDING NUMBERS DAAL03-92-G-0001		
6. AUTHORS Vincent McKoy				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) California Institute of Technology Pasadena, California 91125		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER AR029534.1-EG-SOI		
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words)  The objective of our work is to harness the computational power of large parallel computers to obtain electron collision cross sections needed in modelling the chemical and physical properties of flowfields over hardbodies. Such electron collision data are very fragmentary, often not available, and difficult to measure.  We have developed a scalable implementation of our formulation of electron-molecule collisions. With these scalable algorithms we routinely achieve aggregate performance in the tens of gigaflop range on available computer system. This capability has allowed us to carry out sophisticated first-principles calculations of electron collisions with species such as N <sub>2</sub> , O <sub>2</sub> , CO, NO, and H <sub>2</sub> O.				
14. SUBJECT TERMS		20001122 021		15. NUMBER OF PAGES
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

ENCLOSURE 1

DTIC QUALITY INSPECTED 4

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-1  
298-102

*Electron-Molecule Collisions  
and  
UV Signatures in Flowfields*

*Final Technical Report  
September 2000*

*U.S. Army Research Office  
Contract Number: DAAL03-92-G-0001*

*Performance Period: December 1991 November 1998*

*Principal Investigator: Vincent McKoy  
E-mail: mckoy@its.caltech.edu*

*Institution: California Institute of Technology  
Pasadena, CA 91125*

## *I. Background and Objectives*

Electrons produced in the bow shock of missiles can play an important role in determining the chemical and physical properties of hypersonic flowfields. Due to non-equilibrium overshoots, electron temperatures can rise above 10,000 K in such flowfields. Furthermore, as flight velocities increase, electron number densities and collisions become more significant. However, electron collision data needed in simulations of these flowfields are very fragmentary and often not available. This is particularly true for near-threshold excitation processes where the incident electrons barely have enough energy to drive the excitation and for collisions with metastable states and with fragments such as OH.

Our strategy for obtaining these electron-collision cross section data, particularly those that are experimentally inaccessible, has been to calculate them. Calculations of electron-collision cross sections at the low energies of interest in flowfields, are also difficult. In fact, although the physical principles in low-energy electron-molecule collisions are well understood, and although several methods have been developed for numerical studies of these collisions, progress in their application has been very limited. What is required is not just a method and algorithm by which these cross sections can be calculated in principle, but an implementation that makes relevant problems feasible. The thrust of our effort has hence been to develop innovative scalable implementations of our formulation of electron-molecule collisions<sup>1,2</sup> with which we could harness the computational power of larger parallel computers to obtain electron-collision data needed in modeling flows.

## *II. Summary of Progress and Accomplishments*

At the low energies of interest in these flowfields, an accurate quantum-mechanical treatment of the full  $3(N+1)$ -dimensional space of the Schrödinger's equation that governs the collision of an electron with an  $N$ -electron molecule is necessary; low-order approximations, such as the Born

approximation, are not applicable. Furthermore, because direct numerical integration of Schrödinger's equation in this  $3(N+1)$ -dimensional space is not practical, most recent studies rely on the use of variational approximations to obtain the scattering amplitudes for these collisions. Our studies are based on the Schwinger multichannel (SMC) method,<sup>1</sup> an extension of Schwinger's variational principle. This SMC variational principle was specifically developed for applications to low-energy electron-molecule collisions and possesses compelling computational advantages. However, these come at the cost of evaluating a Green's operator whose matrix representation is obtainable only by numerical quadrature of a class of integrals.<sup>2</sup> Indeed, evaluating and transforming the data for this quadrature is the principal computational task in applications. Not surprisingly, this task had been a bottleneck and hindrance to wider applications of the SMC on sequential computers such as the Cray Y-MP.

We have successfully developed efficient, scalable algorithms for carrying out the main computational tasks in these calculations on large distributed-memory parallel computers consisting of several hundreds of interconnected microprocessors. Our strategy for "parallelization" of these two main tasks — namely, evaluation of a large number (typically  $10^{12}$  to  $10^{15}$ ) of elementary integrals and their subsequent transformation to obtain the physical matrix elements — is as follows. The first of these tasks is perfectly parallel, meaning that on a multiprocessor machine different elementary integrals can be evaluated simultaneously on different processors with effectively zero parallel overhead. Although the second step — transformation of integrals stored in the microprocessors — involves communication and synchronization overheads, it can be formulated in terms of multiplication of large, distributed matrices, a highly efficient procedure on parallel computers. With these scalable algorithms we routinely achieve aggregate performance in the tens of gigaflops range. This capability, which to our knowledge is unmatched, has allowed us to carry out sophisticated first-principles calculations of electron collisions with species such as  $N_2$ ,  $CO$ ,  $NO$ ,  $H_2O$ ,  $OH$ , and  $CO_2$ .

Highlights of these application include

- In a joint effort with the experimental group of Professor H. Ehrhardt of Kaiserslautern University in Germany, we have completed a comprehensive study of the cross sections for electron-impact excitation of the low-lying excited states of CO from threshold to about 4 eV above threshold. These studies should provide the most reliable estimates of the cross sections for excitation of CO in the technically important, but experimentally challenging, near-threshold region. The availability of both measured and calculated differential and integral cross sections offer a unique opportunity for an in-depth comparison of the measured and calculated cross sections. Cross sections were also calculated for impact energies well beyond the threshold region.

- We have also carried out a combined computational – experimental study of the cross section for electron-impact excitation at the low-lying excited states of N<sub>2</sub>. Differential and integral cross sections were calculated for excitation of seven low-lying states (A  $^3\Sigma_u^+$ , B  $^3\Pi_g$ , W  $^3\Delta_u$ , a  $^1\pi_g$ , w  $^1\Delta_u$ , a'  $^1\Sigma_u^-$ , and B'  $^3\Sigma_u^-$ ). Critical comparisons of these calculated cross sections with recent measurements by S. Trajmar of the Jet Propulsion Laboratory were also carried out for the A  $^3\Sigma_u^+$ , B  $^3\Pi_g$ , W  $^3\Delta_u$ , and a  $^1\Pi_g$  states. These studies have provided the first robust values of the cross sections for excitation of the metastable A  $^3\Sigma_u^+$  state of N<sub>2</sub> at near-threshold electron impact energies.

- We have studied the cross sections for near-threshold excitation of the lowest-lying excited state of OH by low-energy electrons. OH fragments are produced in flows of interest via dissociation of water vapor. No measured data are available for these cross sections.

- Cross sections for electron-impact excitation of H<sub>2</sub>O to low-lying excited electronic states, which subsequently fragment to H and OH, were calculated.

## References

1. K. Takatsuka and V. McKoy, Phys. Rev. A **30**, 1734 (1994)
2. C. Winstead, P.G. Hipes, M.A.P. Lima, J. Chem. Phys. **94**, 5455 (1991)
3. C. Winstead and V. McKoy, Adv. At. Mol. Opt. Phys. **36**, 183 (1996)

### III. *The results of this research appear in the following publications.*

1. Collisions of Low-Energy Electrons with CO<sub>2</sub>  
C.-H. Lee, C. Winstead, and V. McKoy  
J. Chem. Phys. **111**, 5056 (1999)
2. Absolute Differential Cross Sections for Electron-Impact Excitation of CO Near Threshold  
J. Zobel, U. Mayer, K. Jung, H. Ehrhardt, H. Pritchard, C. Winstead and V. McKoy  
J. Phys. B. **29**, 839 (1996)
3. Electron Collisions with Nitrous Oxide  
C. Winstead and V. McKoy  
Phys. Rev. A **57**, 3589 (1998)
4. Electron-NO Collisions  
F.J. Paixao, M.A.P. Lima, and V. McKoy  
Phys. Rev. A **53**, 1407 (1996)
5. Highly Parallel Computational Techniques for Electron-Molecule Collisions  
C. Winstead and V. McKoy  
Advances in Atomic, Molecular, and Optical Physics **36**, 183 (1996)
6. Electron Scattering by Small Molecules  
C. Winstead and V. McKoy  
Advances in Chemical Physics **96**, 103 (1996)

7. Parallel Computation of Electron-Molecule Collisions  
C. Winstead, H. Pritchard, and V. McKoy  
Computational Science and Engineering **2**, 34 (1995)
8. Electron Scattering by Water Molecules  
L.E. Machado, M.-T. Lee, L.M. Brescansin, M.A.P. Lima and  
V. McKoy  
J. Phys. B **28** 467 (1995)
9. Studies of Electron-Molecule Collisions on Massively Parallel Computers  
C. Winstead and V. McKoy  
Modern Electronic Structure Theory (World Scientific, 1995)
10. Electronic Excitation of CH<sub>4</sub> by Low-Energy Electron Impact  
C. Winstead, Q. Sun, V. McKoy, J.L.S. Lima, and M.A.P. Lima  
J. Chem. Phys. **98**, 2130 (1993)
11. Electron-Impact Excitation of [1.1.1] Propellane  
C. Winstead, Q. Sun, and V. McKoy  
J. Chem. Phys. **97**, 9483 (1992)
12. Electronic Excitation of Carbon Monoxide by Low-Energy Electron Im-  
pact  
Q. Sun, C. Winstead, and V. McKoy  
Phys. Rev. A1, **46**, 6987 46 (1992)
13. Low-Energy Electron Scattering by C<sub>3</sub>H<sub>6</sub> Isomers  
C. Winstead, Q. Sun, and V. McKoy  
J. Chem. Phys. **96**, 4246 (1992)
14. Studies of Electron Collisions with Polyatomic Molecules Using  
Distributed-Memory Computers  
C. Winstead, P.G. Hipes, M.A.P. Lima, and V. McKoy  
J. Chem. Phys. **94** 5455 (1991)

IV. *Scientific personnel supported by this project:*

Principal Investigator: Vincent McKoy, Professor of Theoretical  
Chemistry

Senior Scientist: Carl Winstead

Graduate Student: Chuo-Han Lee

Honors: Distinguished Lecture Series, Louisiana State University and  
Texas A & M Univeristy.